

SUPPLEMENTAL FIGURES

Sammels et al. (2010) – “Polycystin-2 activation by inositol 1,4,5-trisphosphate-induced Ca^{2+} release requires its direct association with the inositol 1,4,5-trisphosphate receptor in a signaling microdomain”




FIGURE S1. No effect of IP_3 on the interaction between TRPP2 and the IP_3R . GST-pull down using GST or GST-LBD and TRPP2-CT-HIS in the presence of 0, 1 or 10 μM IP_3 . The upper panel shows a Ponceau-red staining of the input GST-fusion proteins. The lower immunoblot shows the staining for the HIS-fusion protein TRPP2-CT-HIS.



FIGURE S2. Sequence of the C-terminus of TRPP2. Alignment of the sequence of the C-terminus of TRPP2 in human, mouse, rat, chicken, zebrafish and polycystin-2 like1 (PKD2L1 or TRPP3) or -2 (PKD2L2 or TRPP5) is shown (made in CLC Sequence Viewer 6). Several domains are indicated on the human TRPP2 sequence: EF-hand (1,2) (orange), the ER-retention signal (3) (blue), the acidic cluster (green), S⁸¹², a CK2 phosphorylation site (4) (encircled) and the coiled-coil region (2,5) (purple).



A**B**

FIGURE S3. Adenoviral expression of wild-type TRPP2. TRPP2^{-/-} cells were transduced with the indicated titers of an adenovirus for wild-type TRPP2. *A*, an immunoblot with staining for TRPP2. *B*, confocal pictures of the cells with a red staining for F-actin and a green staining for the myc-tagged TRPP2. All images were made confocally with the same settings.




FIGURE S4. Adenoviral expression of TRPP2 mutants. TRPP2^{-/-} cells were transduced with a control adenovirus or a TRPP2 adenovirus to re-introduce wild-type TRPP2 or the mutants D⁵⁰⁹V and R⁷⁴⁰X. **A**, 15 µg lysate of transduced TRPP2^{-/-} cells (lane 1-2-3) or LLC-PK1 cells endogenously expressing TRPP2 (lane 4) or over-expressing TRPP2 (obtained by plasmid over-expression) (lane 5) was analysed by SDS-PAGE. An immunoblot is shown with staining for TRPP2 using an antibody directed against the common N-terminal sequence of TRPP2. **B**, confocal pictures of the TRPP2^{-/-} cells with a red staining for F-actin and a green staining for TRPP2. All images were made confocally and with the same settings. Background signal in cells treated with a control virus is shown in the upper left panel.




FIGURE S5. Effect of TRPP2 on a submaximal and maximal agonist-induced Ca^{2+} release in intact cells. TRPP2^{-/-} cells were treated with either control virus (black) (titer 1/1000) or a TRPP2 virus to re-introduce wild-type TRPP2 (red) (titer 1/1000). The ratio of emitted fluorescence of Fura2 was monitored and 10 and 100 μ M ATP were added as indicated in the presence of 3 mM EGTA. $[Ca^{2+}]_{cyt}$ was derived after *in situ* calibration. Panel A shows the averaged traces of at least three independent experiments. Panel B shows the increase in $[Ca^{2+}]_{cyt}$ (nM) after agonist addition, averaged for at least three independent experiments. Panel C shows the quantification of the area under the curve in A.U. Results are shown as means \pm S.E.M.. Statistically significant differences are labelled with * for $p < 0.05$, using a Student t-test (paired two-tailed).




FIGURE S6. ER loading in permeabilized cells transduced with different adenovirus preparations. A, Ca^{2+} uptake by the ER in TRPP2^{-/-} cells transduced with a control virus (black) (titer 1/1000) or a TRPP2 virus to re-introduce wild-type TRPP2 (red) (titer 1/1000) or the mutants D⁵⁰⁹V (green) (titer 1/704) and R⁷⁴⁰X (green) (titer 1/1124). Ca^{2+} uptake in permeabilized cells was initiated by ATP addition. Results are expressed as F/F_0 of MagFluo4 fluorescence (means \pm S.E.M. for three independent experiments). B, maximal $F/F_0 \pm$ S.E.M. for the four conditions.




FIGURE S7. Effect of TRPP2 on TG-induced Ca^{2+} release in intact cells. TRPP2^{-/-} cells were treated with either a control virus (black) (titer 1/1000) or a TRPP2 virus (red) (titer 1/1000) to re-introduce wild-type TRPP2. The ratio of emitted fluorescence of Fura2 was monitored and 10 μM TG was added as indicated in the presence of 3 mM EGTA. Results are shown as mean $[\text{Ca}^{2+}]_{\text{cyt}}$ (as F_{340}/F_{380}) \pm S.E.M. for four independent experiments.

REFERENCES

1. Schumann, F., Hoffmeister, H., Bader, R., Schmidt, M., Witzgall, R., and Kalbitzer, H. R. (2009) *J. Biol. Chem.* **284**, 24372-24383
2. Celic, A., Petri, E. T., Demeler, B., Ehrlich, B. E., and Boggon, T. J. (2008) *J. Biol. Chem.* **283**, 28305-28312
3. Cai, Y., Maeda, Y., Cedzich, A., Torres, V. E., Wu, G., Hayashi, T., Mochizuki, T., Park, J. H., Witzgall, R., and Somlo, S. (1999) *J. Biol. Chem.* **274**, 28557-28565
4. Cai, Y., Anyatonwu, G., Okuhara, D., Lee, K. B., Yu, Z., Onoe, T., Mei, C. L., Qian, Q., Geng, L., Witzgall, R., Ehrlich, B. E., and Somlo, S. (2004) *J. Biol. Chem.* **279**, 19987-19995
5. Yu, Y., Ulbrich, M. H., Li, M. H., Buraei, Z., Chen, X. Z., Ong, A. C., Tong, L., Isacoff, E. Y., and Yang, J. (2009) *Proc. Natl. Acad. Sci. U. S. A.* **106**, 11558-11563